

Examination: Continuous Optimization

3TU- and LNMB-course, Utrecht December 22, 2009, 13.00-16.00

Ex. 1

- 2 (a) Given $a \in \mathbb{R}^n$, show that the matrix aa^T is positive semidefinite.
- 3 (b) Show that a symmetric matrix $A \in \mathbb{R}^{n \times n}$ is positive semidefinite if and only if $A \bullet C \geq 0$ holds for all semidefinite matrices C .
(Here, for symmetric matrices, $A \bullet C$ denotes the "inner product", $A \bullet C = \sum_{i,j} a_{ij}c_{ij}$)

Ex. 2 Consider the convex problem

$$(CO) \quad \min f(x) \quad \text{s.t.} \quad g_j(x) \leq 0, \quad j = 1, \dots, m, \quad x \in \mathbb{R}^n$$

with convex functions $f, g_j \in C^1(\mathbb{R}^n, \mathbb{R})$. Suppose a feasible point \bar{x} satisfies the KKT-conditions (Karush-Kuhn-Tucker) with a multiplier vector $\bar{y} \geq 0$.

- 3 (a) Show that (\bar{x}, \bar{y}) is a saddle point for the Lagrangian function $L(x, y)$ of (CO).
- 3 (b) Show also that (\bar{x}, \bar{y}) is a solution of the Wolfe-Dual (WD) of (CO).

Ex. 3 Let $f_i : C \rightarrow \mathbb{R}, i \in I := \{1, \dots, m\}$ be convex functions on the convex compact set $C \subset \mathbb{R}^n$. Define for $x \in C$ the function f by:

$$f(x) = \min \left\{ \sum_{i=1}^m \lambda_i f_i(x_i) \mid x = \sum_{i=1}^m \lambda_i x_i, \sum_{i=1}^m \lambda_i = 1; x_i \in C, \lambda_i \geq 0 \forall i \in I \right\}$$

Show that $f(x)$ is the greatest convex function $g(x)$ such that $g(x) \leq f_i(x) \forall x \in C$ and $\forall i \in I$, in the following way:

- 3 (a) With the epigraphs $\text{epi}(f_i)$ we consider the set $F := \text{conv} \{ \text{epi}(f_i), i = 1, \dots, m \}$. Show now that $F = \text{epi}(f)$ holds and conclude that f is convex on C .
(By definition, $\text{conv} \{ \text{epi}(f_i), i = 1, \dots, m \}$ is the set

$$\left\{ \sum_{i=1}^m \lambda_i z_i \mid \sum_{i=1}^m \lambda_i = 1; \lambda_i \geq 0, z_i \in \text{epi}(f_i), i \in I \right\},$$

i.e. the set F is the smallest convex set containing all sets $\text{epi}(f_i), i = 1, \dots, m$.)

- 3 (b) Show that $f(x) \leq f_i(x) \forall x \in C$ and $\forall i \in I$ holds and that f is the greatest convex function with this property (i.e., for all convex functions $g(x)$ such that $g(x) \leq f_i(x) \forall x \in C$ and $\forall i \in I$, we have $g(x) \leq f(x), \forall x \in C$).

$$\begin{aligned} & (x^T a a^T x) \\ & (a^T x)^2 > 0 \end{aligned}$$

$$\begin{aligned} & x^T a a^T x \rightarrow \\ & a a^T = x x^T \\ & \downarrow \\ & \|a\|^2 \cdot \|x\|^2 \geq 0 \\ & x^T a a^T x \geq 0 \end{aligned}$$

Ex.4 We consider the unconstrained minimization problem: $\min_{x \in \mathbb{R}^n} f(x)$ with $f \in C^1(\mathbb{R}^n, \mathbb{R})$.

2 (a) Let H be a positive definite (real) $n \times n$ matrix and let $\nabla f(x_k) \neq 0$. Show that $d_k = -H\nabla f(x_k)$ is a descent direction for f in x_k .

3 (b) Given a (non-positive definite) symmetric (real) $n \times n$ matrix A , show that there is a number σ_0 such that for all $\sigma > \sigma_0$ the matrices $A + \sigma I$ are positive definite. (Here, I is the unit matrix). *Hint: Consider the number $\min_{\|x\|=1} x^T A x$*

Ex. 5 Consider the unconstrained minimization problem with the quadratic function $q(x) = \frac{1}{2}x^T A x + b^T x$, where A is a positive definite matrix.

Determine the (global) minimizer \bar{x} of q and show that for any starting point x_0 the Newton iteration computes the minimizer \bar{x} in one step.

Ex. 6 Consider the problem (in connection with the design of a cylindrical can with height h , radius r and volume at least 2π such that the total surface area is minimal):

$$(P) : \min f(h, r) := 2\pi(r^2 + rh) \quad \text{s.t.} \quad -\pi r^2 h \leq -2\pi, \quad (\text{and } h > 0, r > 0)$$

4 (a) Compute a (the) solution (\bar{h}, \bar{r}) of the KKT conditions of (P). Show that (P) is not a convex optimization problem.

2 (b) Show that the solution (\bar{h}, \bar{r}) in (a) is a local minimizer. Why is it the unique global solution?

Hint: Use the sufficient optimality conditions

Ex. 7 We consider the constrained program

$$(P) \quad \min f(x) \quad \text{s.t.} \quad x \in \mathcal{F} \quad \text{where} \quad \mathcal{F} := \{x \in \mathbb{R}^n \mid g_j(x) \leq 0, j \in J\}$$

with $f, g_j \in C^1(\mathbb{R}^n, \mathbb{R})$ and $J = \{1, \dots, m\}$.

Let d_k be a strictly feasible descent direction in $x_k \in \mathcal{F}$, i.e.,

$$\nabla f(x_k)^T d_k < 0, \quad \nabla g_j(x_k)^T d_k < 0, \quad \forall j \in J_{x_k}$$

holds. Show that for any $t > 0$, t small enough, we have:

$$f(x_k + td_k) < f(x_k) \quad \text{and} \quad x_k + td_k \in \mathcal{F}$$

Points: 36+4=40

1	a : 2	2	a : 3	3	a : 3	4	a : 2	5	: 4	6	a : 4
	b : 3		b : 3		b : 3		b : 3				b : 2
7	: 4										

A copy of the lecture-sheets may be used during the examination.

Good luck!